



Incidence of Obesity and Malnutrition Among School-Aged Children of The District Malakand

Usman Saeed², Dr. Umair Islam³, Muhammad Ilyas⁴, Dr. Eidul Ahad⁵, Iqbal Khan Rahman⁶, Muhammad Yaseen¹, Noor Muhammad⁷, Iqbal Muhammad⁷, Hamza Khan¹, Abbas Khan¹, Shakir Ullah¹, Said Ullah¹

¹Department of Pathology Takht Bhai Institute of Health and Management Sciences Mardan, Pakistan

²Department of Zoology GC University Lahore

³Department of Internal Medicine Khyber Teaching Hospital Peshawar Pakistan.

⁴Department of Pathology Riphah International University Malakand Campus, Pakistan.

⁵Department of Dentistry Spinghar Medical College Pakistan

⁶Department of Microbiology University of Swat, Pakistan.

⁷Department of Microbiology Kohat University of Science & Technology (KUST), Kohat, Pakistan.

*Correspondin shakirullah1992@gmail.com +923469058726

ABSTRACT

Obesity and malnutrition are worldwide health problem so the aims of the study were to assist the obesity and nutritional deficiencies among school-aged children. Institutional based cross-sectional study designs were used. In which total of 6000 children between the ages of 4-15 years were studied. A systematic random sampling technique was applied for sample collection. A total of 6000 students were participated in this study. The numbers of male students and females students were same. All the students were from primary and high school. The overall results show that out of 6000 students the ratio of underweight were 415(7%) and obese (6.1%) respectively. Our study shows that nutritional factors are important since obesity and overweight increase with socioeconomic status as well as with nutritional intake. So Obesity was found High socioeconomic status groups should be directed for overweight while underweight and nutritional deficiency is a problem of lower socioeconomic status. Meat, Milk intake and lack of physical activity have been identified in the current study. We observed that a strong relation was found in nutrition with food intake and socioeconomic status.

Keywords: Obesity, Overweight, Underweight, Malnutrition

Received: 1-5-2024

Accepted: 14 -5-2024

Published: 1-6-1024

INTRODUCTION

Obesity is one of the most prevalent nutritional disease of children and adolescents in many developed and developing countries. The World Health Organisation (**WHO 1998**) has declared overweight as one of the top ten health risks in the world and one of the top five in developed

nations. Existing (**WHO 2002**) standards and data from 79 developing countries including a number of industrialized countries suggest that about 22 million children five years old are overweight worldwide. (**WHO 1998**) Once considered a problem of affluence, obesity is fast growing in many developing countries also. (**WHO 2000**) Even in countries like India, which are typically known for high prevalence of under nutrition, a significant proportion of overweight and obese children now coexist with those who are under nourished. (**Popkin et al 2001**) Increasing relative weight trends in populations have caused much concern among health care providers. Limited studies have been conducted on the prevalence of overweight (OW) and obesity (OB) in the children belonging different socio economic groups in the National Capital Territory (NCT) of Delhi hence the present study was conducted.

Obesity has emerged as one of the global health problems with 200 million school-aged children world-wide categorized as being overweight/obese, of which 40-50 million are obese (**Shikhakhandelwal et al 2016**). The factors attributing to increasing childhood obesity are increased intake of high-calorie foods that are low in vitamins, minerals and micronutrients coupled with decreased physical activity (**Kaushik et al 2011**). Various studies done in India from 2002-2012 indicate a rising trend in the prevalence of overweight and obesity in children and adolescents (**Shah et al 2011**).

MATERIALS AND METHODS

Study area:

A cross-sectional study was conducted in the local area of district Malakand. All the primary, middle and high schools were enlisted in this study. Socio economic were made including Lower Income socio economic Group (LIG), Middle socio economic Income Group (MIG) and High Income socio economic Group (HIG). All the Government and private schools were measured for obesity.

Sample size

This study was undertaken for the Exploration of vitamin D deficiency in school aged children of the local area of District Malakand. We retrospectively studied the records of a total of 6000 children aged between 4-15 June 2023 to December 2023. All patients were subjected to a careful physical examination. Weights were measured using a calibrated digital scale. Height measurements were done in triplicate to the nearest millimeter using a calibrated stadiometer. Body mass index (BMI) were calculated according to the formula [weight (kg)/height (m)²]. Patients with a history of a chronic disorder or on any medication that may alter vitamin D metabolism were excluded from the study. Serum calcium (Ca), phosphorus (P), magnesium (Mg), alkaline phosphatase (ALP) and glucose levels were measured using the enzymatic colorimetric method (Roche Integra 800), while serum 25 hydroxyl vitamin D [25(OH) D] levels were measured by high-performance liquid chromatography (Shimadzu UFLC).

Demographics information:	General information	Dietary Habits
Child's Name: _____	Do you have breakfast every day before going to school? (Yes/No)	What is your favorite healthy food?
Age: _____ Gender: _____	How many meals do you typically eat in a day?	How often do you drink water in a day?
Grade/Class: _____	Do you eat fruits and vegetables daily? (Yes/No)	Less than 3 glasses 3-5 glasses 6-8 glasses More than 8 glasses
School Name: _____	How often do you consume fast food? (Yes/No)	Are you aware of the importance of a balanced diet? (Yes/No)
Family background:	Anthropometric Measurements	Do you receive any nutrition education at school? (Yes/No)
How many people live in the child's household.?	Height (cm): __	
Male and female ratio in child's household.?	Weight (kg): __	
Family income status?	BMI (Body Mass Index): __	
Hereditary diseases in family?		

According to WHO equation: $BMI = \frac{\text{Weight (kg)}}{\text{Height (m)}^2}$

S.no	Status	Student BMI	Normal BMI	HB level g/dl
1	Anemia	20	18.5-24.9	>12,>14
2	Underweight	18	18.5-24.9	<12,14
3	Malnutrition	17.3	18.5-24.9	<12,14
4	Stunting	11.2	18.5-24.9	<12,14
5	Wasting	9.5	18.5-24.9	<12,14

HB: hemoglobin

Study Area and Period

The study was conducted in local area of district Malakand, which is located in Khyber Pakhtunkhwa.

Study Design

Institutional based cross-sectional study designs were used.

Study Population

All secondary and primary school students (age group wise the source population, whereas sampled or selected students were the study population of this study.

Sample Size Determination

A Total of 6000 children between the ages of 4-15 years were studied. A systematic random sampling technique was applied for sample collection.

Clinical Examination

Routine blood tests

This is done to assess anemia and other vitamin and mineral deficiencies. There may be dehydration, low blood sugar and signs of severe infection as is evident by raised white blood cell counts.

Blood tests in children

Routine blood tests in children include those for blood glucose, blood counts, urine for routine examination.

Levels of iron in blood, folic acid and vitamin B 12 are also done. For protein estimation other tests including

Normal ranges of ferritin 10 to 150 ng/mL for children 4th years to 14 years.

MCV normal range is 80 to 95 for children

Analysis & Interpretation Analysis and Interpretation of data

Data will analyze and interpret by using M word, Origin8 and Excel. Frequency and Percentage were calculated for all quantitative variables.

Ethical Consideration

The subjects were briefed about the study.

- Consent was taken from the subjects after explaining the purpose of study for the collection of data.
- **Questionnaire form**

Demographics information:	General information	Dietary Habits
Child's Name:	Do you have breakfast every day before going to school? (Yes/No)	What is your favorite healthy food?
Age: Gender:	How many meals do you typically eat in a day?	How often do you drink water in a day?
Grade/Class:	Do you eat fruits and vegetables daily? (Yes/No)	Less than 3 glasses
School Name: _	How often do you consume fast food? (Yes/No)	3-5 glasses
Family background:	Anthropometric Measurements	6-8 glasses
How many people live in the child's household.?	Height (cm):	More than 8 glasses
Male and female ratio in child's household.?		Are you aware of the importance of a balanced diet? (Yes/No)

Family income status?	Weight (kg):	Do you receive any nutrition
Hereditary diseases in family?	BMI (Body Mass Index):	education at school? (Yes/No)

S.no	Serum 25(OH)D	Status
1	Serum 25(OH)D	Deficient < 50nmol/L
		Insufficient 50-75nmol/L
		Sufficient <75nmol/L
2	Mean Serum 25(OH)D	55± 6nmol/L

Statistical Analysis

Statistical analysis was performed by using Origin8 and MS office word 2010.

RESULTS AND DISCUSSION

Total of 6000 students were participated in this study. The numbers of male students and females students were same. All the students were from primary and high school. The overall results shows that out of 6000 students the ration of underweight were 415(7%) and obese (6.1%) respectively (Table.1).

Table.1 Overall nutritional assessment of school-aged children

Socioeconomic status	Male			Female		
	Normal	Underweight	Obese	Normal	Underweight	Obese
Low Income	850	150(2.5%)	00(0%)	900	100(1.66%)	0(0%)
Middle income	900	50(0.83%)	50(0.83%)	850	90(1.5%)	60(1%)
High Income	830	10(0.16%)	160(2.66%)	885	15(0.25%)	100(1.66%)
Total	2580	210(3.5%)	210(3.5%)	2635	205(3.41%)	160(2.66%)

Our study shows that nutritional factors are important since obesity and overweight increase with socioeconomic status as well as with nutritional intake. So Obesity was found High socioeconomic status groups should be directed for overweight while underweight and nutritional deficiency is a problem of lower socioeconomic status. Meat, Milk intake and lack of Physical activity is some of the other factors that have been identified in the current study that are shown in table.1 and 2.

Table.2 Nutritional intake wise assessment of school children

MEAL TIMINGS				
Do You Take	Always	Sometimes	Rarely	Never
Breakfast	135	200	250	120
Lunch	165	135	120	180
Tea time	50	150	130	220
Dinner	250	115	100	80
Food Response and Reactions				
Cravings	Aversions		Unsuitability	
250	180		170	
WATER INTAKE (1Glass=250ml)				

Our study also shows those nutritional intakes are important meanwhile obesity and overweight increase with socioeconomic status as well. So Obesity was found High socioeconomic status groups should be directed for overweight while underweight and nutritional deficiency is a problem of lower socioeconomic status. Meat, Milk intake and lack of physical activity are some of the other factors that have been identified in the current study that are shown in table.1 and 2.




Figure.1 socioeconomic status wise nutritional assessment of school-aged children

Gender wise nutritional assessment of the current study shows that the underweight and obese were found high in male group than the female with the prevalence rate of male underweight

210(3.5% and obese 210(3.5%) while female were found slightly low underweight 205(3.41%) and obese 160(2.66%) respectively has been shown in figure no.2.




Figure.2 Gender wise nutritional assessment of school-aged children

DISCUSSION

Our study shows that nutritional factors are important since obesity and overweight increase with socioeconomic status as well as with nutritional intake. So Obesity was found High socioeconomic status groups should be directed for overweight while underweight and nutritional deficiency is a problem of lower socioeconomic status. Meat, Milk intake and lack of physical activity are some of the other factors that have been identified in the current study. Similarly a study also conducted by (Chhatwal et al 2004) and (Khader et al 2009) in a population, 70% of all obese children belonged to the higher SEC, while of the underweight children, 63.3% were in the lower SEC. This finding is consistent with the view of previous studies that obesity in developing countries increases with socioeconomic class. Meat consumption was high across the board; however, it was significantly higher in children from higher SES. Within overweight and obese kids, there was not a single report of children not eating meat at all. This was however, not true of many underweight children belonging to lower SEC, who reported not having meat at all. Meat intake has been found to be associated with obesity and its cardiovascular complications. Children in Switzerland showed a direct association between intake of meat products and overweight.

Gender wise nutritional assessment of the current study shows that the underweight and obese were found high in male group than the female with the prevalence rate of male underweight 210(3.5% and obese 210(3.5%) while female were found slightly low underweight 205(3.41%) and obese 160(2.66%) respectively. Same study were also performed by (Wijga et al 2010) in which the total 2,131 children, 19.4% were overweight (18.8% of boys and 19.9% of girls) and 5.6% were obese (5.6% of boys and 5.5% of girls). Watching television [2 h/day, daily pocket money [20 piasters (1 piaster = 1.42 cents), having overweight or obese mother/father were significantly associated with increased odds of both overweight and obesity. Age C 10 years, female gender, and family size of B4 were significantly associated with being overweight and total monthly family income [300 Jordanian Dinars (JDs), (1 JD = \$1.42) was associated with obesity.

CONCLUSION

We observed that a strong relation was found in nutrition with food intake and socioeconomic status. The time expended in sleeping was a progressive factor for keeping a balance between weight and height. There was also a nutritional association found between male and female children.

ACKNOWLEDGEMENTS

We appreciate the support of the Department of Pathology THQ Takht Bhai towards the success of the study. We will also like to thank the staffs of the School teachers and school heads for their technical support.

ETHICAL CONSIDERATIONS

Participation was voluntary and confidentiality was assured to all respondents. They were informed about the study's objectives and that the data collected would be used only for the stated research purposes.

AVAILABILITY OF DATA AND MATERIALS

The study materials are available from the corresponding author upon request.

REFERENCES

- Chatterjee P. India sees parallel rise in malnutrition and obesity. Lancet. 2002;** 360:1948.
Shah, C., Diwan, J., Rao, P., Bhabhor, M., Gokhle, P., & Mehta, H. (2008). Assessment of obesity in school children. *Calicut Medical Journal*, 6(3), e2.
- Chhatwal J, Verma M, Riar SK (2004)** Obesity among pre-adolescent and adolescents of a developing country (India). *Asia Pac J Clin Nutr* 13: 231–235.
- Jagadesan, S., Harish, R., Miranda, P., Unnikrishnan, R., Anjana, R. M., & Mohan, V. (2014).** Prevalence of overweight and obesity among school children and adolescents in Chennai. *Indian pediatrics*, 51, 544-549.
- Kaushik JS, Narang M, Parakh A. Fast food consumption in children. Indian Pediatr. 2011;**48:97-101.

- Khader, Y., Irshaidat, O., Khasawneh, M., Amarin, Z., Alomari, M., & Batieha, A. (2009). Overweight and obesity among school children in Jordan: prevalence and associated factors. *Maternal and child health journal*, 13, 424-431.
- Spinelli, A., Buoncristiano, M., Kovacs, V. A., Yngve, A., Spiroski, I., Obreja, G., ... & Breda, J. (2019). Prevalence of severe obesity among primary school children in 21 European countries. *Obesity facts*, 12(2),244-258.
- WHO consultation on obesity. Special issues in the management of obesity in childhood and adolescence. In: obesity preventing and managing the global epidemic Geneva: WHO, 1998; pp 231-47.
- WHO (World Health Organisation). 2002. The World Health Report: Reducing Risks, Promoting Healthy Life. Geneva: World Health Organisation.
- WHO (World Health Organisation). 1998a. Obesity: Preventing and Managing the global Epidemic. Report of WHO Consultation on Obesity. WHO, Geneva.
- WHO (World Health Organisation), International Association for the Study of Obesity (IASO) and International Obesity task Force (IOTF). 2000. The Asia-Pacific Perspective: Redefining Obesity and its treatment. Geneva: World Health Organisation.
5. Popkin BM., D Horton, S Kim, A Mahal and J Shuigao. Trends in diet nutritional status and diet related non communicable diseases in China and India: The economic costs of the nutritional transition, *Nutr Rev.* 2001;59:379-90. Shikhakhandelwal, D. P. M., & Gupta, M. L. Prevalence of obesity and overweight in private schools of Jaipur city in 11-17 year age group children.
- Wijga, A. H., Scholtens, S., Bemelmans, W. J., de Jongste, J. C., Kerkhof, M., Schipper, M., ... & Smit, H. A. (2010). Comorbidities of obesity in school children: a cross-sectional study in the PIAMA birth cohort. *BMCPublicHealth*, 10,1-10