



Nutritional status of schoolchildren with recurrent tonsillitis in Takht Bhai District, Mardan, Pakistan

Usman Shah¹, Dr Umair Islam², Usman Saeed³, Ismail Shahid⁴, Muhammad Iqbal Khan Rahman⁵, Maryam Bibi⁶, Basit Ali⁷, Shakir Ullah⁶, Yaseen Khan⁸, Muhammad Kashif Gul⁸

¹Department of Health Khyber Medical University Peshawar, Paksitan 25000

²Department of Internal Medicine KTH Peshawar Pakistan, 25000

³Department of Zoology GC University Lahore Pakistan, 54000

¹Department Public Health Health services academy Islamabad Pakistan, 44000

⁵Department of Microbiology University of Swat, 45000

⁶Department of Microbiology Abasyn University Peshawar Pakistan, 25000

⁷Department of Veterinary Sindh Agriculture University Tando Jam Pakistan, 65081

⁸Department of Chemistry Abdul Wali Khan University Mardan, 23500

*Corresponding Author: shakirullah1992@gmail.com

+923469058726

ORCID: <https://orcid.org/0000-0003-1731-9344>

ABSTRACT

The recurrent tonsillitis can be the significance of weaknesses in immune system, the particular etiology of recurrent tonsillitis is not clear. Therefore aims of the study are to explore Nutritional status among school children with recurrent tonsillitis in Takht Bhai District Mardan Pakistan of aged 4 to 15 years. This observational, cross sectional study was conducted in Takht Bhai District Mardan from March 2023 to October 2023. In the current research work a total of 250 samples were isolated from school-going children. In which 75(30%) were found positive for nutritional deficiency. Among the all samples nutritional anemia was found more frequent 30(12%), leading by malnutrition 25(10%), moderate prevalence were shows by underweight 10(4%) and stunting 8(3.2%) prevalence rate. while very low frequency was shows by wasting(2(0.8%) respectively. Out of 250 children the number of male was slightly higher than females. Nutritional Anemia and malnutrition were observed in the study population respectively. The prevalence of nutritional Anemia among the study population varied with age. Children of the 12–15 years group had a significantly high prevalence of nutritional Anemia compared with the lower age group. Although the prevalence of nutritional anemia was higher in both age groups and sexes, it was not significant. Nutritional Anemia prevalence was significantly higher among the school going children. The difference between the age groups as well as nutritional status remained significant as risk factors of nutritional anemia also the main relation. Over all ratio of underweight and stunting were found moderate among all ages and gender wise distribution of nutritional status. This study revealed that there were weak relations found between nutritional status and tonsillitis

Keywords: Anemia, BMI, Nutritional status, Malnutrition, under nutrition, Tonsillitis

INTRODUCTION

Variety of poor outcomes including growth retardation, historically, the science of nutrition developed in and development of psychosocial difficulties. Water low 1995 of disease entities brought about reported that etiology of linear growth retardation is by inadequate diet (Sireesha & Rajani, 2015). Nutritional status is the condition of multi-factorial but has been explained by three major health of an individual as influenced by nutrient intake factors: poor nutrition, high levels of infection and utilization in the body (De Onis, 2017).

In developing world, approximately nutrients in preparation for rapid growth of adolescence 146 million children are underweight, (Handa, Ahamad, Kesari, & Prasad, 2008). Therefore, it becomes very important to know the nutritional status of school going children; the building machine was used to measure the body weight to the blocks of state and country and hence the present nearest 0.5 kg (Best, Neufingerl, Van Geel, van den Briel, & Osendarp, 2010). Malnutrition is a major public health problem in Pakistan, a South Asian nation with over 130 million people. Half of its children aged five years or less are stunted, over a third (38%) are underweight, and a quarter of all births are low birth weight (Best et al., 2010).

These high levels of malnutrition contribute to about half of the 740,000 child deaths that occur every year in Pakistan (Shivaprakash & Joseph, 2014). In view of the scale of the problem in children under five, nutritional programmes in Pakistan during the last few decades have been targeted at this age group. However, malnutrition is a significant problem in older children as well, a fact that is often overlooked by policy makers and programme managers (Fazili, 2012). Though little is known about the state of nutrition in this older group, studies conducted in the 1980s indicate that malnutrition is a significant problem in this population, with prevalences ranging from 47-70% in male school children in rural Pakistan (Karak, Maiti, Das, & Karmakar, 2018; Rezaeian et al., 2014).

The situation among school-aged children in urban squatter settlements in Pakistan is even less well known. These settlements contain a large proportion of the rapidly growing urban population, with high levels of malnutrition already documented in the under-five child population (Eze, Oguonu, Ojinnaka, & Ibe, 2017). To assess the nutritional status of 7-10 year old Anthropometric. Nutritional assessment is a depth evaluation of an individual food, nutrient intake lifestyle and their medical history. Nutritional assessment of collecting information in order to make decisions about the nature and cause of nutritional related health issues that can affect an individual.

We can evaluate the nutritional status of an individual by using these components whether, he or she is obese overweight or underweight. Because obesity and overweight may lead to chronic diseases such as diabetes, heart failure, hypertension and so on. While under nutrition can lead to osteoporosis getting frequent ill, teeth, skin and hair problems, iron deficiency anemia, malnutrition, pre-mature birth. In 2015, the study on teachers indicate that 34% were overweight, 16% were anemic and 2% had thin-built. According to WHO the prevalence of obesity has been doubled and 2.8 million individuals died due to obesity and 15.8 billion were considered obese. (khushnuma amin, Oct 2015)

Another study was conducted on females of Pakistan showed that approx 60% of the population consumed less than recommended daily allowance and 40% consumed less than 80% of the recommended daily allowance. In Pakistan 19.9% lactating females consumed calories less than 70% of the recommended dietary allowance while 54% a pregnant female in Pakistan were anemic as compare to India, were more than 50% of the populations were anemic, while iodine deficiency also has significance effects on poor pregnancy outcomes.

A high mortality rate in Pakistan is due to under nutrition and almost 25 to 30% of babies under 25,00g at the birth. And low birth weight is a significant reason behind high mortality and this high mortality rate was due to poor availability and accessibility of food. Workload and pressure of house-hold work directly effects on the nutritional status of females. (Chatterjee & Lambert, 1989).

A study was conducted on depression, anxiety and stress level of female school teachers according to the study teaching not only affect physically but also mentally because a lot of energy is used in the class rooms, in family commitments, which is a source of stress and depression..in Egypt primary teachers take 24classes per week, while secondary level teachers took 18classesper week. The study indicated that 61.3%had teaching experience more than 10years while 91%teachers were not satisfied with their pay scale. The prevalence of severe, moderate and mild depression were (0.7%, 2.8%and 19.7%),while the prevalence of extremely severe, moderate and mild anxiety level was (19.7%, 7.0% and 23.2%). These results showed that there is need to bring interventions to overcome such issues and order to prevent teachers from increasing psychological issues which would affect mentally and physically on nutritional status of teachers (Allam, 2017).

Showed that the prevalence of anaemia and undernutrition are each associated with significant morbidity and mortality, with higher rates among children, particularly in sub-Saharan Africa. Anaemia is a condition where due to low blood haemoglobin concentration (Kateera et al 2015) the oxygen-carrying capacity of red cells is insufficient to meet the body's physiologic needs. This condition affects individuals globally and has significant adverse health consequences, as well as adverse impacts on social and economic development (Alcázar 2013).Childhood anaemia is considered a severe public health problem in Sub-Saharan Africa (62.5%) and in Cameroon in particular where a prevalence of 63.2% was reported in 2011.

Stunting is associated with impaired cognitive development, reduced academic achievement, and

decreased physical work capacity in adulthood, with a negative cost on the economic development of societies. While the global stunting prevalence fell from 39.6 to 23.8% between 1990 and 2014, the scenario is quite different in Africa, with an increase . Nevertheless, in some localities in the Mount Cameroon area, the prevalence of stunting fell from 49.9% to 17.1% (Sumbele et al 2015). The impact of nutritional status on malaria may differ due to the heterogeneity of the population under study, species of the parasite, and other factors involved in the host and parasite relationship.

AMIS OF THE STUDY

To explore Nutritional status among school children with recurrent tonsillitis in Takht Bh District Mardan Pakistan of aged 4 to 15 years.

MATERIALS AND METHODS

This observational, cross sectional study was conducted in Takht Bhai District Mardan from March 2023 to October 2023. After ethical approval, students of 4-15 years were selected from the Govt schools located in this area. Demographic information along with weight and height of the selected children were taken and plotted to gender specific, growth charts.

The School Health Program was carried out on regular basis in the Rural Block. For the purpose of the present paper, the survey was findings from schools surveyed from March 2023 to October 2023 was included. The age of the children was determined using school records. In the schools nutritional status of children were assessed as follows: Weight: Measured using a floor type weighing scale with due respect to the standardization of the equipment and procedure. The measurements were taken to the nearest .5Kg. Height: well taken using a measuring tape applied to the wall. The measurements were taken with children barefoot with their back of heels, buttocks and head touching the wall. Readings well be taken to the nearest

5cm.WHOz- score system were used to classify the nutritional status of children.

Study Area and Period

The study was conducted in Takht Bhai Mardan, which is located Khyber Pakhtunkhwa.

Study Design

Institutional based cross-sectional study design were used.

Study Population

All secondary and primary school students (age group wise the source population, whereas

sampled or selected students were the study population of this study.

Sample Size Determination

A Total of 250 children between the ages of 4-15 years were studied. A systematic random sampling technique was applied for sample collection.

Lab investigations and Tests: All patients were subjected to:

Complete Blood count and reticulocyte count. Patients with microcytosis underwent the following:

• Serum Iron and ferritin

An automatic hematological analyzer and Biochemistry analyzer were used for clinical examination

Routine blood tests

This is done to assess anemia and other vitamin and mineral deficiencies. There may be dehydration, low blood sugar and signs of severe infection as is evident by raised white blood cell counts.

Diagnosis of malnutrition in children

In children weight and height is measured and compared with the charts showing the expected average height and weight for a child of that age. Some children are persistently smaller for age and may be genetically so.

Blood tests in children

Routine blood tests in children include those for blood glucose, blood counts, urine for routine examination.

Levels of iron in blood, folic acid and vitamin B 12 are also done. For protein estimation other tests including

Normal ranges of ferritin 10 to 150 ng/mL for children 4th years to 14 years.

MCV normal range is 80 to 95 for children

ANALYSIS & INTERPRETATION

► Analysis and Interpretation of data

Data will analyze and interpret by using M word, Origin 16 and Excel. Frequency and Percentage were calculated for all quantitative variables.

5. ETHICAL CONSIDERATION

- The subjects were briefed about the study.
- Consent were taken from the subjects after explaining the purpose of study for the collection of data.

Questionnaire form

| | | |
|--|---|---|
| Demographics information: Child's Name: _____ Age: _____ Gender: _____ | General information Do you have breakfast every day before going to school? (Yes/No) How many meals do you typically | Dietary Habits What is your favorite healthy food? How often do you drink water in a |
|--|---|---|

| | | |
|---|--|--|
| Grade/Class: _____ School Name: _____ Family background: How many people live in the child's household? Male and female ratio in child's household? Family income status? Hereditary diseases in family? | eat in a day? Do you eat fruits and vegetables daily? (Yes/No) How often do you consume fast food? (Yes/No) Anthropometric Measurements Height (cm): ____ Weight (kg): ____ BMI (Body Mass Index): ____ | day? Less than 3 glasses 3-5 glasses 6-8 glasses More than 8 glasses Are you aware of the importance of a balanced diet? (Yes/No) Do you receive any nutrition education at school? (Yes/No) |
|---|--|--|

| | | | | |
|--|------------------|------------------|----------------------|--------------|
| FOOD FREQUENCY QUESTIONNAIRE FOR NUTRITIONAL STATUS OF SCHOOLCHILDREN WITH RECURRENT TONSILLITIS IN TAKHT BHAI DISTRICT, MARDAN, PAKISTAN | | | | |
| FOOD FREQUENCY QUESTIONNAIRE (FFQ) | | | | |
| Code | | | | |
| MEAL TIMINGS | | | | |
| Do You Take | Always | Sometimes | Rarely | Never |
| Breakfast | 100 | 50 | 40 | 60 |
| Lunch | 150 | 55 | 30 | 15 |
| Tea time | 120 | 35 | 50 | 45 |
| Dinner | 200 | 10 | 25 | 15 |
| FOOD RESPONSE AND REACTIONS | | | | |
| Cravings | Aversions | | Unsuitability | |
| | | | | |
| WATER INTAKE (1Glass=250ml) | | | | |

According to WHO formula $BMI = \frac{\text{weight}}{\text{Height}^2(M^2)}$

| s.no | status | Student BMI | Normal BMI | HB level |
|------|--------------|-------------|------------|-------------|
| 1 | Anemia | 20BMI | 18.5-24.9 | >12,>14g/dL |
| 2 | Underweight | 18BMI | 18.5-24.9 | <12,14g/dL |
| 3 | Malnutrition | 17.3BMI | 18.5-24.9 | <12,14g/dL |
| 4 | Stunting | 11.2BMI | 18.5-24.9 | <12,14g/dL |
| 5 | wasting | 9.5BMI | 18.5-24.9 | <12,14g/dL |

RESULTS

In the current research work a total of 250 samples were isolated from school-going children. In which 75(30%) were found positive. This study revealed that there were weak relations found between nutritional status and tonsillitis. The prevalence of tonsillitis in anemic children was 7(2.8%), malnutrition 3(1.2%), Underweight 2(0.8%), stunting 1(0.4%) while 0% in wasting. "Table 2" Among the all samples nutritional anemia was found more frequent 30(12%), leading by malnutrition 25(10%), moderate prevalence were shows by underweight 10(4%) and stunting 8(3.2%) prevalence rate. while very low frequency was

shows by wasting(2(0.8%) respectively. Out of 250 children the number of male was slightly higher than females. The prevalence of nutritional Anemia among the study population varied with age. Children of the 12–15 years group had a significantly high prevalence of nutritional Anemia compared with the lower age group “Table 2”. Although the prevalence of nutritional anemia was higher in both age groups and sexes, it was not significant. The distribution of malnutrition by nutritional severity is summarized in “Table 1”. Anemia prevalence was significantly higher among the school going children. Over all ratio of underweight and stunting were found moderate among all ages and gender wise distribution of nutritional status.

Table.1 Class wise and gender wise nutritional status of school-going

| S.no | Class | Anemia | | Malnutrition | | Underweight | | Stunting | | Wasting | |
|------|-------------------|--------|---|--------------|---|-------------|---|----------|---|---------|---|
| | | M | F | M | F | M | F | M | F | M | F |
| 1 | 1 st J | 4 | 3 | 3 | 2 | 2 | 1 | 0 | 1 | 1 | 0 |
| 2 | 1 st S | 3 | 3 | 2 | 3 | 1 | 1 | 1 | 0 | 1 | 0 |
| 3 | 2 nd | 3 | 2 | 2 | 1 | 0 | 0 | 1 | 0 | 0 | 0 |
| 4 | 3 rd | 2 | 2 | 0 | 1 | 0 | 1 | 0 | 1 | 0 | 0 |
| 5 | 4 th | 1 | 2 | 1 | 0 | 1 | 0 | 1 | 1 | 0 | 0 |
| 6 | 5 th | 1 | 1 | 2 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| 7 | 6 th | 1 | 1 | 2 | 2 | 1 | 1 | 0 | 0 | 0 | 0 |
| 8 | 7 th | 1 | 0 | 1 | 1 | 1 | 0 | 0 | 1 | 0 | 0 |
| 9 | 8 th | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 10 | 9 th | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 11 | 10 th | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |




Figure.1 Over all nutritional status of school going children

This study revealed that there were weak relations found between nutritional status and tonsillitis.

The prevalence of tonsillitis in anemic children was 7(2.8%), malnutrition 3(1.2%), Underweight 2(0.8%), stunting 1(0.4%) while 0% in wasting. “Table 2” The prevalence of malnutrition and stunting varied significantly within the age group with the lower age group having higher prevalence. More males were anemic 16 (6.4%), malnourished 14 (5.6%), underweight 6 (2.4%), stunted 4 (1.6%) when compared with females nutritional anemia 14 (5.6%), malnutrition 11 (4.4%), underweight 4 (1.6), stunting 4 (1.6%) while wasting with zero prevalence rate 0 (0%) respectively though the difference was not significant in malnutrition and stunting. The difference in prevalence of underweight among the sexes was significant with the males having a higher prevalence. Bivariate analysis revealed children of the 0–5 year age group were significantly at odds of being malnourished “Table 3”

Table.2 Gender wise nutritional status of school going children

| S.no | Nutritional status | Male | | Female | |
|------|--------------------|--------------------|-------------|--------------------|-------------|
| | | Nutritional status | Tonsillitis | Nutritional status | Tonsillitis |
| 1 | Anemia | 16 (6.4%) | 3(1.2%) | 14 (5.6%) | 4(1.6%) |
| 2 | Malnutrition | 14 (5.6%) | 2(0.8%) | 11 (4.4%) | 1(0.4%) |
| 3 | Underweight | 6 (2.4) | 1(0.4%) | 4 (1.6) | 1(0.4%) |
| 4 | Stunting | 4 (1.6%) | 0(0%) | 4 (1.6%) | 1(0.4%) |
| 5 | Wasting | 2 (0.8%) | 0(0%) | 0 (0%) | 0(0%) |
| 6 | -- | 42(17%) | 6(2.4%) | 33(13%) | 7(2.8%) |




Figure.2 Over all nutritional status of school going children




Figure.3 Over all nutritional status of school going children

Age-wise nutritional status in males shows that the high prevalence rates were found between

4-7 years of children. Nutritional anemia 12 (4.8%) led by Malnutrition 10 (4%) less prevalence was seen in Underweight 5 (2%) Stunting 5 (2%) while some prevalence was seen for wasting 2 (0.8%). Between the age group 8-11 the prevalence ratio was nutritional anemia 10 (4%) led by Malnutrition 8 (3.2%) less prevalence was seen in Underweight 3 (1.2%) and Stunting 2 (0.8%) while no prevalence was seen for wasting. Also, high prevalence was seen in nutritional anemia and malnutrition 8 (3.2%), 7 (2.8%) very low prevalence rates were seen between the 12-15 age group in underweight, Stunting, and wasting Table.4

Table.3 Age wise nutritional status of school going children

| Ages | Anemia | Malnutrition | Underweight | Stunting | Wasting |
|-------|------------------|-----------------|-----------------|-----------------|----------------|
| 4-7 | 12 (4.8%) | 10 (4%) | 5 (2%) | 5 (2%) | 2 (0.8) |
| 8-11 | 10 (4%) | 8 (3.2%) | 3 (1.2%) | 2 (0.8%) | 0 (0%) |
| 12-15 | 8 (3.2%) | 7 (2.8%) | 2 (0.8%) | 1 (0.4%) | 0 (0%) |
| Total | 30(12%) | 25(10%) | 10(4%) | 8(3.2%) | 2(0.8%) |

Age-wise nutritional status in males shows that the high prevalence rates were found between 4-7 years of children. Nutritional anemia 8 (3.2%) led by Malnutrition 6 (2.4%) less prevalence was seen in Underweight 4 (1.6%) Stunting 2 (0.8%) while some prevalence was seen for wasting 2 (0.8%). Between the age group 8-11 the prevalence ratio was nutritional anemia 5 (2%) leading by Malnutrition 4 (1.6%) less prevalence was seen in Underweight 2 (0.8%) and Stunting 2 (0.8%) while no prevalence was seen for wasting. Very low prevalence rates were seen between the 12-15 age group Table.5

Table.4 Age wise nutritional status of male school going children

| Ages | Anemia | Malnutrition | Underweight | Stunting | Wasting |
|-------|-----------------|-----------------|-----------------|-----------------|-----------------|
| 4-7 | 8 (3.2%) | 6 (2.4%) | 4 (1.6%) | 2 (0.8%) | 2 (0.8%) |
| 8-11 | 5 (2%) | 4 (1.6%) | 2 (0.8%) | 2 (0.8) | 0 (0%) |
| 12-15 | 3 (1.2%) | 4 (1.6%) | 2 (0.8%) | 0 (0%) | 0 (0%) |
| Total | 16(6.4%) | 14(5.6%) | 8(3.2%) | 4(1.6) | 2(0.8%) |

Age-wise nutritional status in males shows that the high prevalence rates were found between 4-7 years of children. Nutritional anemia 8 (3.2%) led by Malnutrition 6 (2.4%) less prevalence was seen in Underweight 4 (1.6%) Stunting 2 (0.8%) while some prevalence was seen for wasting 2 (0.8%). Between the age group 8-11 the prevalence ratio was nutritional anemia 5 (2%) led by Malnutrition 4 (1.6%) less prevalence was seen in Underweight 2 (0.8%) and Stunting 2 (0.8%) while no prevalence was seen for wasting. Very low prevalence rates were seen between the 12-15 age groups Table.5

Table.5 Age wise nutritional status of female school going children

| Ages | Anemia | Malnutrition | Underweight | Stunting | Wasting |
|-------|-----------------|-----------------|-----------------|-----------------|---------------|
| 4-7 | 7 (2.8%) | 5 (2%) | 2 (0.8%) | 3 (1.2%) | 0 (0%) |
| 8-11 | 5 (2%) | 4 (1.6%) | 2 (0.8%) | 1 (0.4%) | 0 (0%) |
| 12-15 | 2 (0.8%) | 2 (0.8%) | 2 (0.8%) | 0 (0%) | 0 (0%) |
| Total | 14(5.6%) | 11(4.4%) | 6(2.4%) | 4(1.6%) | 0(0%) |

DISCUSSION

In the current research work, a total of 250 samples were isolated from school-going children. The samples were collected from male and female students of age 4-15 years. Among the all samples nutritional anemia was found more frequent 30(12%), led by malnutrition at 25(10%), moderate prevalence was shown by underweight 10(4%), and stunting at 8(3.2%) prevalence rate. while very low frequency was shows by wasting(2(0.8%) respectively. Out of 250 children, the number of males was slightly higher than females. Nutritional Anemia and malnutrition were observed in the study population respectively. The prevalence of nutritional Anemia among the study population varied with age. Children in the 12–15 years group had a significantly higher prevalence of nutritional Anemia compared with the lower age group. (**Ahsan et al 2020**) also conducted a study in which a total of 571 children, 348 (56.4%) were boys and 223 (43.6%) were girls. Nearly 89 (15.5%) children including 52 (10.5%) boys and 37(6.4%) girls had normal anthropometric indices. Stunting was the most frequent anthropometric failure (n=219, 38.3%) followed by wasting (n=163, 28.51%) and underweight (n=100, 17.5%) respectively. Gender disparity was observed in the distribution of malnutrition with boys having a higher frequency of stunting, wasting, and underweight than girls.

In the current study the tonsillitis status revealed that there were weak relations found between nutritional status and tonsillitis. The prevalence of tonsillitis in anemic children was 7(2.8%), malnutrition 3(1.2%), Underweight 2(0.8%), stunting 1(0.4%) while 0% in wasting. (**Metgudmath et al 2023**) also conducted a study in which 84% of the patients had a iron deficiency anaemia, 56% had low serum iron, There was a significant corelation between anaemia and low serum iron among the study population, the Spearman's correlation coefficient r value was 0.824 and the p value was less than 0.05 between iron deficiency anaemia and recurrent tonsillitis with higher grades of hypertrophy and symptoms as well.

Although the prevalence of nutritional anemia was higher in both age groups and sexes, it was not significant. Nutritional Anemia prevalence was significantly higher among school-going children. The difference between the age groups as well as nutritional status remained significant as risk factors of anemia were also the main relation. The ratio of underweight and stunting was found moderate among all ages and gender-wise distribution of nutritional status. The prevalence of malnutrition and stunting varied significantly within the age group with the lower age group having higher prevalence. More males were anemic 16 (6.4%), malnourished 14 (5.6%), underweight 6 (2.4%), stunted 4 (1.6%) when compared with females anemia 14 (5.6%), malnutrition 11 (4.4%), underweight 4 (1.6), stunting 4 (1.6%) while wasting with zero prevalence rate 0 (0%) respectively though the difference was not significant in malnutrition and stunting. The difference in prevalence of underweight among the sexes was significant with the males having a higher prevalence. Bivariate analysis revealed children of the 0–5 year age group were significantly at odds of being malnourished. Similar work was also performed by (**Shakir Ullah et al 2023**). Conducted a study in which out of 1500 school-going students 600(40%) were found positive and 900(60%) were found normal according to age. Gender-wise analysis shows that in male students 210(35%) were found positive for stunting and 150 (25%) male students were underweight. In overall female students, 135(22.5%) were found stunting and 105 (17.5%) were found underweight. According to age group wise between 4 to 10 years 90 (10%) female students were found stunting and 60(6.6%) were found underweight. While in the age of 11 to 15 years 48 (5.3%) students were stunted and 42 (4.6%) students were underweight. The difference in stunted and underweight showed more boys than girls. In the pre-nursery group, more boys were stunted than girls, the ratio being 70%:30%, on the contrary, underweight was more in boys than in girls, with a ratio of 57%:43%. In the

Primary section, both stunting and underweight were more in boys as compared to girls. The ratios were 54.8%:45.2% and 82.9%:17% respectively.

Age-wise nutritional status in males shows that the high prevalence rates were found between 4-7 years of children. Nutritional anemia 12 (4.8%) led by Malnutrition 10 (4%) less prevalence was seen in Underweight 5 (2%) Stunting 5 (2%) while some prevalence was seen for wasting 2 (0.8%). Between the age group 8-11 the prevalence ratio was nutritional anemia 10 (4%) led by Malnutrition 8 (3.2%) less prevalence was seen in Underweight 3 (1.2%) Stunting 2 (0.8%) while no prevalence was seen for wasting. Also, high prevalence was seen in nutritional anemia and malnutrition 8 (3.2%), 7 (2.8%) very low prevalence rates were seen between the 12-15 age group in underweight, Stunting, and wasting. While a study also conducted by (Farhin et al 2021). The mean age of the sample was 9.38 ± 4.14 with the maximum number of children (49.1%) in the age bracket of 5-9. Out of 1710 children, 54.4% had normal weight for age, 25.3% were underweight, 7.5% overweight and 12.8% were found to be obese. Stunting was found to be 26%. The prevalence of being underweight was higher than overweight /obesity, particularly in younger and higher age groups as indicated by a p-value of 0.000. Compared with females, male students had a significantly higher frequency of being underweight and stunted as reflected by p-values of 0.004 and 0.000 respectively. Univariate analysis also showed a strong association between age and nutritional status as mean weight increased from 39.22 ± 5.21 to 63.50 ± 4.66 and height from 35.67 ± 5.76 to 113.73 ± 29.22 with advancing age.

CONCLUSION

Nutritional Anemia prevalence was significantly higher among the school going children. The difference between the age groups as well as nutritional status remained significant as risk factors of nutritional anemia also the main relation. Over all ratio of underweight and stunting were found moderate among all ages and gender wise distribution of nutritional status. This study revealed that there were weak relations found between nutritional status and tonsillitis

RECOMMENDATION

The present study has successfully documented the nutritional status in terms of the prevalence of underweight, stunting and Tonsillitis among the rural school-going children of Tehsil Takht Bhai district Mardan. Nutritional Anemia prevalence was significantly higher among the school going children. The difference between the age groups as well as nutritional status remained significant as risk factors of nutritional anemia also the main relation. Over all ratio of underweight and stunting were found moderate among all ages and gender wise distribution of nutritional status. This study revealed that there were weak relations found between nutritional status and tonsillitis. So some other research works also needed to find the relation and treatment of the recurrent tonsillitis in school aged children.

Acknowledgement

Authors feel grateful to the Microbiology Department of Abasyn University peshawar and Takht Bhai Institute of Health and Management Sciences for providing facilities during research activities.

Competing Interests

The authors declare that they have no competing interests.

Authors' Contributions

All authors equally contributed in the designing, experiments and wrote the manuscript. All authors read and approved the final manuscript.

REFERENCES

- Adnan M S Fakir, M Wasiquir Rahman Khan (2015) Determinants of malnutrition among urban slum children in Bangladesh. *Health Econ Rev* 5(1): 59. thinness among Pakistani primary school children. *Biomed Central*. 2011;11:790-BMC .Public health.K. Park. Parks Textbook of Preventive and Social Medicine. 18th ed. Jabalpur: Banarsidas
- Ahmed, I.F. and A.H. Gilani, 1998. Effect of protein energy malnutrition on serum protein levels in school boys. *Pak. J. Med. Res.*, 27: 192-5.
- Ahsan, S., Saleh, Z., Sheikh, S. A., Fahim, M. F., Memon, M. S., & Shakil, S. (2020). Nutritional status of school going children of 5-15 years of age: urban slums scenario in Karachi, Pakistan. *Biostatistics and Biometrics Open Access Journal*, 10(2), 22-26.
- Ahsan, S., Saleh, Z., Sheikh, S. A., Fahim, M. F., Memon, M. S., & Shakil, S. (2020). Nutritional status of school going children of 5-15 years of age: urban slums scenario in Karachi, Pakistan. *Biostatistics and Biometrics Open Access Journal*, 10(2), 22-26.
- Akor, F. (2010). Nutritional status of newly enrolled primary school. *Pakistan journal of nutrition*, 9(12), 1166- 1170.
- Alcázar L. The economic impact of anaemia in Peru. In: 2013: MISC; 2013.
- Njunda AL, Ngouadjeu DT, Nsagha D, Nyanjoh EM, Kwenti T, Assob NJ. Haematological profile of children with malaria in Kumba Health District, South West Region Cameroon. *Afr J Integrated Health*. 2016;6:23–9
- Akiyama T, Pongvongsa T, Phrommala S, Taniguchi T, Inamine Y, Takeuchi R, et al. Asymptomatic malaria, growth status, and anaemia among children in Lao People's Democratic Republic: a cross-sectional study. *Malar J*. 2016;15:499.
- Arlappa, N., Balakrishna, N., Laxmaiah, A., & Brahmam, G. N. V. (2010). Relevance of anaemia among rural pre-school children of West Bengal, India. *Annals of human biology*, 37(2), 231-242.
- Batool, S., Shaheen, A., Rehman, R., Qamar, S., Raza, S. A., Jabeen, R., & Nisa, F. (2012). To assess the nutritional status of primary school children in an Urban school of Faisalabad. *Pak J Med Health Sci A*, 4, 160.
- Bernard, H.R., 1995. *Research Methods in Anthropology*. Sage, California, pp: 360- 92..
- Best, C., Neufingerl, N., Van Geel, L., van den Briel, T., & Osendarp, S. (2010). The nutritional status of school-aged children: why should we care? *Food and nutrition bulletin*, 31(3), 400-417.
- Begum, M., & Nessa, Z. (2008). Nutritional status of school going children of a selected school of Dhaka city. *Bangladesh Journal of Scientific and Industrial Research*, 43(1), 97-102.
- Bhanot; 2005.
- International institute of population Sciences. National Family Health Survey-2, 1998-99. Mumbai, India: IIPS 2000. in the first 5 yrs for children in developing countries. *Lancet* 2007, 309; 60-70.
- Bhutta Z.A, 2000 . Why has so little changed in maternal and child health in south Asia? *Br. Med. J.*, 321: 809-812.
- Black RE, Allen LH ,Bhutta ZA et al 2017. Maternal and child undernutrition global and regional exposures and health consequences. *Lancet* 2008, 371; 243-260 Victoria G, Adair L, Fall C et al. Maternal and child undernutrition, consequences for adult health and human capital. *Lancet* 2008, 371; 340-353. Grantham Mcgregor S, Cheung YB, Cento S, Glewer P, Kichter L, Strupp B. Development potential in the first 5 yrs for children in developing countries. *Lancet* 2007, 309; 60-70.
- Boussougou-Sambe ST, Awono-Ambene P, Tasse GC, Etang J, Binyang JA, Nouage LD, et al. Physical integrity and residual bio-efficacy of used LLINs in three cities of the South-West Region of Cameroon 4 years after the first national mass-distribution campaign. *Malar J*. 2017;16:31.

- Brown, J.L. and L.P. Sherman, 1995. Policy implications of new scientific knowledge. *J. Nutr.*, 125 (Suppl 8): 2281S-2284S. Caldwell, J. and P. McDonald, 1982. Influence of maternal education on infant and child mortality: levels and causes. *Health Policy and Education*, 2: 251-267.
- Caulfeld LE, Richard SA, Black RE. Undernutrition s an underlying cause of malaria morbidity and mortality in children less than five years old. *Am J Trop Med Hyg.* 2004;71(suppl 2):55–63.
- Chhabra, P., S. Garg, N. Sharma and R.D. Bansal, 1996. Health and nutritional status of boys aged 6-12 years ina children observation home. *Indian J. Public Health*, 40: 126-9.
- De Onis, M. (2017). Child growth and development. *Nutrition and health in a developing world*,
- Eyong EEJ, Kengne-Ouafo AJ, Chounna PW, Datchoua-Poutcheu FR, Wanji S. Altitudinal variation in the parasitological and entomological indices of malaria **around Mount Cameroon, South West Region of Cameroon.** *J Parasitol Vector Biol.* 2016;8:74–85.
- Farhin, S., Jaffry, T., Zafar, S., & Rashid, F. (2021). Frequency and assessment of Nutritional Status of school going children in rural areas of Islamabad. *Pakistan Journal of Medical Sciences*, 37(5), 1475.
- Eze, J., Oguonu, T., Ojinnaka, N., & Ibe, B. (2017). Physical growth and nutritional status assessment of school children in Enugu, Nigeria. *Nigerian journal of clinical practice*, 20(1), 64-70.
- Fazili, A. (2012). Nutritional status of school age children (5-14 years) in a rural health block of North India (Kashmir) using WHO Z-score system. *Online Journal of Health and Allied Sciences*, 11(2).
- Fazili, A., Mir, A. A., Pandit, B. M., Bhat, I. A., Rohul, J., & Shamila, H. (2012). Nutritional status of school age children (5-14 years) in a rural health block of North India (Kashmir) using WHO Z-score system. *Online Journal of Health and Allied Sciences*, 11(2 (2)).
- Fellows E (2019) 10 Facts About Poverty in Karachi.
- UN-Habitat, 2003 UN-Habitat (2003) The challenge of slums - global report on human settlements 2003. London, UK: United Nations Human Settlements Programme.
- Hadju, V., K. Abadi, L.S. Stephenson, N.N. Noor and H.O. Mohammed, 1995. Intestinal helminthiasis, nutritional status, and their relationship; a cross-sectional study in urban slum school children in Indonesia. *Southeast Asian J. Tropical Med. and Public Health*, 26: 719-29.
- Handa, R., Ahamad, F., Kesari, K. K., & Prasad, R. (2008). Assessment of nutritional status of 7-10 years school going children of Allahabad district: A review. *Middle East J Sci Res*, 3(3), 109-115.
- Kateera F, Ingabire CM, Hakizimana E, Kalinda P, Mens PF, Grobusch MP, et al. Malaria, anaemia and under-nutrition: three frequently coexisting conditions among preschool children in rural Rwanda. *Malar J.* 2015;14:440.
- Karak, P., Maiti, R., Das, P., & Karmakar, A. (2018). Assessment of nutritional status of school children in rural and urban areas of Bankura, West Bengal. *Int J Pharm Sci Res*, 9(1), 338-345.
- Kennedy, E. and C. Davis, 1998. US department of agriculture school breakfast program. *Am. J. Clin. Nutr.*, 67: 798S-803S. Martorell, R., 1995. Results and implications of the INCAP followup study. *J. Nutr.*, 125 (Suppl 4):1127S-1138S.
- Kovalskys, I., Rausch Herscovici, C., & De Gregorio, M. (2011). Nutritional status of school-aged children of Buenos Aires, Argentina: data using three references. *Journal of public health*, 33(3), 403-411.
- Mwanik i, E., & Makokha, A. (2013). Nutrition status and associated factors among children in public primary schools in Dagoretti, Nairobi, Kenya. *African health sciences*, 13(1), 38-46.

- Masyithah Darlan, D., Alexandra, T., & Tala, Z. (2017). Soil Transmitted Helminth Infections in Medan: a cross-sectional study of the correlation between the infection and nutritional status among elementary school children. *Family Medicine & Primary Care Review*, (2), 98-103.
- Mansur, D. I., Haque, M. K., Sharma, K., Mehta, D. K., & Shakya, R. (2015). A study on nutritional status of rural school going children in Kavre District. *Kathmandu University Medical Journal*, 13(2), 146-151.
- Mwaniki, E., & Makokha, A. (2013). Nutrition status and associated factors among children in public primary schools in Dagoretti, Nairobi, Kenya. *African health sciences*, 13(1), 38-46.
- Metgudmath, V., Metgudmath, V. B., & Vasudevan, C. (2023). Iron deficiency anaemia as a novel immunological indicator of recurrence in chronic tonsillitis. *IP Journal of Otorhinolaryngology and Allied Science*, 6(1), 12-16
- Ngnie-Teta, I., Receveur, O., & Kuate-Defo, B. (2007). Risk factors for moderate to severe anemia among children in Benin and Mali: insights from a multilevel analysis. *Food and Nutrition Bulletin*, 16(1), 1-10.
- Nolan LB (2015) Slum definitions in urban India: implications for the measurement of health inequalities. *Popul Dev Rev* 41(1): 59-84.
- Nyaaba DA, Brown PK, Agbenorhevi JK. Assessing the association between nutritional status and asymptomatic malaria parasitaemia of children under five years in Ghana. *Pyrex J Nutrition Metabolism*. 2017;2:1-14.
- Rezaeian, S., Ahmadzadeh, J., Esmailnasab, N., Veisani, Y., Shayan, M., & Moradi, N. (2014). Assessment of health and nutritional status in children based on school screening programs. *Nutrition Bulletin*, 28(1), 76-89.
- Rao, A., & Parikh, A. (2015). Prevalence of anemia in school age children. *Journal of advanced Medical and Dental Sciences Research*, 3(6), S32.
- Sarkar, A., Arafat, Y., Alam, M., & Sarkar, J. (2020). Study on Dietary Pattern and Nutritional Status of School Going Children in Navaron, Jashore, Bangladesh. *Am J Food Sci Technol*, 8, 70-4.
- Shanti Ghosh, Dheeraj Shah (2004) Nutritional problems in urban slum children. *Indian pediatr* 41(7): 682-696. in the first 5 yrs for children in developing countries. *Lancet* 2007,309;60-70.
- Shivaprakash, N., & Joseph, R. B. (2014). Nutritional status of rural school-going children (6-12 Years) of Mandya District, Karnataka. *International Journal of Scientific Study*, 2(2), 39-43.
- Sireesha, G., & Rajani, N. (2015). Nutritional status of girls through anthropometric and dietary assessment. *IOSR J Dent Med Sci*, 14(11), 73-78.
- Tomar, S. P., Kasar, P. K., & Tiwari, R. (2016). Study of oral hygienic practices and oral health status among school children in Jabalpur, Madhya Pradesh: a cross-sectional study. *Int J Community Med Public Health*, 3(2), 403-7.
- Unger A (2013) Children's health in slum settings. *Archives of disease in childhood* 98(10): 799-805. United Nations Human Settlements Programme (UN-Habitat) (2003) The challenge of slums ;Global report on human settlement.
- Ullah, S., Khan, H., Saeed, U., & Ahmad, K. (2023). Assessment of nutritional status of school-going children in district Mardan at Khyber Pakhtunkhwa in Pakistan. *Egyptian Journal of Nutrition*, 38(4), 12-18.
4. UN-Habitat, 2010 UN-Habitat (2010) State of the cities 2010-11 - cities for all: Bridging the urban divide. UN-Habitat, Nairobi, Kenya. in the first 5 yrs for children in developing countries. *Lancet* 2007,309;60-70.
- Verma, M., Sharma, P., Khanna, P., Srivastava, R., & Sahoo, S. S. (2021). Nutrition status of school children in Punjab, India: Findings from school health surveys. *Journal of Tropical Pediatrics*, 67(1), fmaa068

- WHO. World malaria report 2015. Geneva: World Health Organization;2016.<http://www.who.int/malaria/publications/world-malaria-report2016/report/en/>. Accessed 27 Apr 2018. Sumbele IU, Ning TR, Bopda OS, Nkuo-Akenji T. Variation in malariometric and red cell indices in children in the Mount Cameroon area following enhanced malaria control measures: evidence from a repeated crosssectional study. *Malar J.* 2014;13:334. Sumbele IUN, Sama SO, Kimbi HK, Taiwe GS. Malaria, moderate to severe anaemia, and malarial anaemia in children at presentation to hospital in the Mount Cameroon area: across sectional study. *Anemia.*2016;2016:5725634.
- Wanzira H, Katamba H, Okullo AE, Agaba B, Kasule M, Rubahika D. Factors associated with malaria parasitaemia among children under 5 years in Uganda: a secondary data analysis of the 2014 Malaria Indicator Survey dataset. *Malar J.* 2017;16:191. Sumbele IUN, Kimbi HK, Ndamukong-Nyanga JL, Nweboh M, Anchang-Kimbi JK, Lum E, et al. Malarial anaemia and anaemia severity in apparently healthy primary school children in urban and rural settings in the Mount Cameroon area: cross sectional survey. *PLoS One.* 2015;10:e0123549.
- WHO. World Malaria Report. Geneva: World Health Organization; 2014.http://www.who.int/malaria/publications/world_malaria_report_2014/en/. Accessed 27 Apr 018. Sumbele IUN, Bopda OSM, Kimbi HK, Ning TR, Nkuo-Akenji T. Nutritional status of children in a malaria meso endemic area: cross sectional study on prevalence, intensity, predictors, influence on malaria parasitaemia and anaemia severity. *BMC Public Health.* 2015;15:1099.
- Waterlow, J.C., 1994. Introduction, causes and achievement of regular and irregular breakfast eaters mechanisms of linear growth retardation (stunting). and noneaters. *Annual Report*, 4: 27-30. *Eur. J. Clin. Nutr.*, 48(1): S1-4. 17.
- Gowri, A.R. and H.J. Sangunam, 2005. Assessment of preschool children. *The Ind. J. Nutr. Dietet.*, 38: 182.
- mental and motor abilities of school going children 19. World Health Organization, 2002. Malnutrition with anemia. *Ind. J. Nutr. Dietet.*, 42: 99-105.